


Immunizations and Developmental Milestones for Your Child from Birth Through 6 Years Old

Child's Name _____

Birth Date _____

		Birth	1 MONTH	2 MONTHS	4 MONTHS	6 MONTHS
Recommended Immunizations	Hepatitis B	<input type="radio"/> HepB	<input type="radio"/> HepB ¹			<input type="radio"/> HepB
	Rotavirus			<input type="radio"/> RV	<input type="radio"/> RV	<input type="radio"/> RV
	Diphtheria, Tetanus, Pertussis			<input type="radio"/> DTaP	<input type="radio"/> DTaP	<input type="radio"/> DTaP
	<i>Haemophilus influenzae</i> type b			<input type="radio"/> Hib	<input type="radio"/> Hib	<input type="radio"/> Hib
	Pneumococcal			<input type="radio"/> PCV	<input type="radio"/> PCV	<input type="radio"/> PCV
	Inactivated Poliovirus			<input type="radio"/> IPV	<input type="radio"/> IPV	<input type="radio"/> IPV
	Influenza					<input type="radio"/> Influenza, first dose ² <input type="radio"/> second dose
Milestones*	<p>Milestones should be achieved by the end of the age indicated.</p> <p>Talk to your healthcare provider about age-appropriate milestones if your child was born prematurely.</p>	<input type="radio"/> Recognizes caregiver's voice	<input type="radio"/> Responsive to calming actions when upset <input type="radio"/> Starts to smile <input type="radio"/> Raises head when on tummy	<input type="radio"/> Developing a social smile <input type="radio"/> Raises chest when lying on tummy <input type="radio"/> Follows objects with eyes <input type="radio"/> Makes cooing noises	<input type="radio"/> Grasps and shakes hand toys <input type="radio"/> Laughs out loud <input type="radio"/> Babbles more expressively <input type="radio"/> Holds head steady	<input type="radio"/> Responds to own name <input type="radio"/> Reaches with one hand <input type="radio"/> Recognizes familiar faces <input type="radio"/> Rolls over
Growth	At each well child visit, enter date, length, weight, and percentile information to keep track of your child's progress.	WEIGHT / PERCENTILE	WEIGHT / PERCENTILE	WEIGHT / PERCENTILE	WEIGHT / PERCENTILE	WEIGHT / PERCENTILE
		LENGTH / PERCENTILE	LENGTH / PERCENTILE	LENGTH / PERCENTILE	LENGTH / PERCENTILE	LENGTH / PERCENTILE
		HEAD CIRCUMFERENCE	HEAD CIRCUMFERENCE	HEAD CIRCUMFERENCE	HEAD CIRCUMFERENCE	HEAD CIRCUMFERENCE

continues on back page

 Shaded boxes indicate the vaccine can be given during shown age range.

VISIT DATE

VISIT DATE

VISIT DATE

VISIT DATE

VISIT DATE

¹ The second dose of HepB may be given either at the 1 month or 2 month visit.
² Two doses given at least four weeks apart are recommended for children aged 6 months through 8 years of age who are getting a flu vaccine for the first time.

* Milestones adapted from *Bright Futures: Guidelines for Health Supervision of Infants, Children, and Adolescents* Third Edition, edited by Joseph Hagan, Jr., Judith S. Shaw, and Paula M. Duncan, 2008, Elk Grove Village, IL: American Academy of Pediatrics.



U.S. Department of Health and Human Services
Centers for Disease Control and Prevention



American Academy of Pediatrics




DEDICATED TO THE HEALTH OF ALL CHILDREN™

Immunizations and Developmental Milestones for Your Child from Birth Through 6 Years Old

Child's Name _____

Birth Date _____

		12 MONTHS	15 MONTHS	18 MONTHS	19 23 MONTHS	2-3 YEARS	4-6 YEARS
Recommended Immunizations	Hepatitis B	<input type="radio"/> HepB (Final dose administered between 6 and 18 months)					
	Diphtheria, Tetanus, Pertussis		<input type="radio"/> DTaP				<input type="radio"/> DTaP
	Haemophilus influenzae type b	<input type="radio"/> Hib					
	Pneumococcal	<input type="radio"/> PCV³					
	Inactivated Poliovirus	<input type="radio"/> IPV					<input type="radio"/> IPV
	Influenza	<input type="radio"/> Influenza , first dose ² <input type="radio"/> second dose (if needed)			Age 2 3 <input type="radio"/> <input type="radio"/> Influenza , first dose ² <input type="radio"/> <input type="radio"/> second dose (if needed)		Age 4 5 6 <input type="radio"/> <input type="radio"/> <input type="radio"/> Influenza , first dose ² <input type="radio"/> <input type="radio"/> <input type="radio"/> second dose (if needed)
	Measles, Mumps, Rubella	<input type="radio"/> MMR					<input type="radio"/> MMR
	Varicella	<input type="radio"/> Varicella					<input type="radio"/> Varicella
Hepatitis A	<input type="radio"/> <input type="radio"/> Hep A⁴						
Milestones*	<p>Milestones should be achieved by the end of the age indicated.</p> <p>Talk to your healthcare provider about age-appropriate milestones if your child was born prematurely.</p>	<input type="radio"/> Cries when mother or father leaves <input type="radio"/> Finger-feeds himself <input type="radio"/> Explores objects in many different ways (shaking, banging, throwing, dropping) <input type="radio"/> Says "dada" and "mama"	<input type="radio"/> Drinks from a cup <input type="radio"/> Scribbles <input type="radio"/> Indicates what he wants by pointing	<input type="radio"/> Says at least 6 words with meaning <input type="radio"/> Points to one body part <input type="radio"/> Walks up steps	<input type="radio"/> Completes simple chores when asked <input type="radio"/> Is spontaneous with affection	<input type="radio"/> Points to object or picture when named <input type="radio"/> Uses 2- to 4-word phrases <input type="radio"/> Follows simple instructions <input type="radio"/> Gradually becomes more interested in other children	<input type="radio"/> Balances on one foot <input type="radio"/> Shows school readiness skills <input type="radio"/> Prints some letters and numbers <input type="radio"/> Has mature pencil grasp <input type="radio"/> Has good articulation/language skills
	Growth	<p>At each well child visit, enter date, length, weight, and percentile information to keep track of your child's progress.</p>	_____ WEIGHT / PERCENTILE _____ LENGTH / PERCENTILE _____ HEAD CIRCUMFERENCE	_____ WEIGHT / PERCENTILE _____ LENGTH / PERCENTILE _____ HEAD CIRCUMFERENCE	_____ WEIGHT / PERCENTILE _____ LENGTH / PERCENTILE _____ HEAD CIRCUMFERENCE	_____ WEIGHT / PERCENTILE _____ LENGTH / PERCENTILE _____ HEAD CIRCUMFERENCE	_____ WEIGHT _____ HEIGHT _____ BMI

 Shaded boxes indicate the vaccine can be given during shown age range.

VISIT DATE

VISIT DATE

VISIT DATE

VISIT DATE

VISIT DATE

VISIT DATE

² Two doses given at least four weeks apart are recommended for children aged 6 months through 8 years of age who are getting a flu vaccine for the first time.

³ Children 2 years old and older with certain medical conditions may need a dose of pneumococcal vaccine (PPSV) and meningococcal vaccine (MCV4). See vaccine-specific recommendations at <http://www.cdc.gov/vaccines/pubs/ACIP-list.htm>.

⁴ Two doses of HepA vaccine are needed for lasting protection. The first dose of HepA vaccine should be given between 12 months and 23 months of age. The second dose should be given 6 to 18 months later. HepA vaccination may be given to any child 12 months and older to protect against HepA. Children and adolescents who did not receive the HepA vaccine and are at high-risk, should be vaccinated against HepA.

* Milestones adapted from *Bright Futures: Guidelines for Health Supervision of Infants, Children, and Adolescents* Third Edition, edited by Joseph Hagan, Jr., Judith S. Shaw, and Paula M. Duncan, 2008, Elk Grove Village, IL: American Academy of Pediatrics.



U.S. Department of Health and Human Services
Centers for Disease Control and Prevention



American Academy of Pediatrics
DEDICATED TO THE HEALTH OF ALL CHILDREN™