

# 2010–11 SEASONAL INFLUENZA VACCINE

## Most children 6 months through 8 years of age need **2** doses


Choose from any of the 3 equivalent tools below (text, table, flow chart) to check how many doses are needed for a child 6 months through 8 years of age:

A child 6 months through 8 years of age is recommended to receive **2** doses of 2010–11 seasonal influenza vaccine at least 4 weeks apart

**UNLESS** determined to have received:

- ✓ at least 1 dose of Monovalent **H1N1** vaccine in 2009–10
- AND**
- ✓ at least 1 dose of Trivalent **Seasonal influenza vaccine** before July 2009 **OR** 2 doses in 2009–10

Number of influenza vaccine doses <b>previously received</b>			Number of doses <b>recommended</b> for the 2010–2011 season
<b>SEASONAL</b>		<b>2009–10 H1N1</b>	
July 2009 – June 2010	Before July 1, 2009		
2	<b>OR</b> 1 or more	2 or 1	<b>1</b>
		0 or unknown	<b>2</b>
1, 0, or unknown (No known receipt of any seasonal doses or only 1 dose in 2009-10)	<b>AND</b> 0 or unknown	2, 1, 0, or unknown	administered at least 4 weeks apart



## Persons 9 years and older need **1** dose of 2010-2011 seasonal influenza vaccine